

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 82111258.8

(22) Date of filing: 06.12.82

(51) Int. Cl.³: **C 07 C 93/14, C 07 C 91/30,**
C 07 C 103/44, C 07 D 285/10,
C 07 D 213/74, C 07 D 215/38,
A 61 K 31/135, A 61 K 31/41,
A 61 K 31/445, A 61 K 31/47

(30) Priority: 14.12.81 US 330542

(43) Date of publication of application: 22.06.83
Bulletin 83/25

(84) Designated Contracting States: AT BE CH DE FR GB IT
LI LU NL SE


(71) Applicant: MERCK & CO. INC., 126, East Lincoln Avenue
P.O. Box 2000, Rahway New Jersey 07065 (US)

(72) Inventor: Jensen, Norman P., 5 Woodcrest Drive, New
Providence New Jersey (US)
Inventor: Chang, Michael N., 746 Austin Street, Westfield
New Jersey (US)

(74) Representative: Abitz, Walter, Dr.-Ing. et al, Abitz, Morf,
Gritschneider, Freiherr von Wittgenstein
Postfach 86 01 09, D-8000 München 86 (DE)

(54) Hydroxybenzylaminobenzenes as anti-inflammatory agents.

(57) Hydroxybenzylaminobenzenes of structural formula (I)



are disclosed wherein:

R is

- (a) fluoro;
- (b) methoxy, ethoxy, n-propoxy or i-propoxy;
- (c) methylthio, ethylthio, n-propylthio or i-propylthio;
- (d) -OCH₂-O;
- (e) -COOH; or
- (f) aryloxy;

R₁ is

- (a) unsubstituted or substituted phenyl; or
- (b) unsubstituted or substituted heteroaryl;

n is 1 or 2. These compounds have been prepared from an appropriate hydroxybenzaldehyde and an amine followed by reduction. These compounds are found to be active topical anti-inflammatory agents.

TITLE OF THE INVENTION

HYDROXYBENZYLAMINOBENZENES AS ANTI-INFLAMMATORY AGENTS

BACKGROUND OF THE INVENTION

5 The present invention relates to novel
hydroxybenzylaminobenzenes useful as topical
anti-inflammatory agents. We have found that the
novel compounds are active in both the peritoneal
macrophage assay and the U.V. erythema assay for
10 topical anti-inflammatory agents. However, these
compounds are rapidly inactivated in vivo and devoid
of any systemic effects.

 Recent studies demonstrated that macrophages
participate in the development and progression of
15 chronic inflammatory diseases such as rheumatoid
arthritis. During the progression of inflammatory
conditions, there is generally an appearance and/or
presence of macrophages and lymphocytes, especially
macrophages and polymorphonuclear leukocytes.
20 Macrophages are known to secrete various products in
response to inflammatory stimuli. For example:

- 1) Neutral proteinases - the destructive peptide bond cleaving enzyme which has been shown to be directly involved in rheumatoid cartilage destruction; and
- 5 2) Prostaglandins (PG) (e.g., E_2 and I_2 by mouse peritoneal macrophages) and other arachidonic acid derivatives derived from both the cyclooxygenase and the lipoxygenase pathways.

10 These arachidonic acid oxygenation products have been identified as the critical mediators of various acute inflammatory conditions.

15 Accordingly, pharmacological agents which are capable of inhibiting the formation, the release, or the function of macrophages may also be effective agents in the treatment of rheumatoid arthritis, emphysema, bronchial inflammation, osteoarthritis, psoriasis, acute respiratory distress syndrome, spondylitis, lupus, gout and other inflammatory

20 diseases.

 With respect to the U.V. erythema assay, it has been shown previously that the U.V. erythema condition is partially the result of a local release of prostaglandins derived oxidatively from

25 arachidonic acid by the action of PG synthetases, e.g., cyclooxygenase and lipoxygenase. Therefore, pharmacological agents which inhibit the erythema are generally considered to be active topical anti-inflammatory agents.

30 Furthermore, anti-inflammatory agents which are not systemically active are advantageous in the sense that they are not subject to the adverse

effects, e.g., gastrointestinal ulcerations and bleeding that often plagued users of systemic NSAIDs (non-steroidal anti-inflammatory agents). Accordingly, an object of this invention is to provide
5 novel hydroxybenzylaminobenzenes as topical anti-inflammatory agents useful in the treatment of dermal inflammatory conditions and pruritus such as sunburn, erythema, eczema, contact dermatitis, and allergic dermatitis, and psoriasis.

10 Another object of this invention is to provide appropriate processes for the preparation of the subject novel compounds.

Still a further object of the present invention is to provide a pharmaceutically acceptable
15 composition containing an effective amount of the active compound for the treatment of various dermatological inflammatory conditions.

Finally, it is the ultimate object of this invention to develop a method of treating dermal
20 inflammation via the administration of an effective amount of the novel compounds as well as a pharmaceutically acceptable composition thereof to a mammalian species in need of such treatment.

25 DETAILED DESCRIPTION OF THE INVENTION

This invention relates to novel derivatives of hydroxybenzylamine of the structural formula (I):

or a pharmaceutically acceptable salt thereof.

Wherein R is

- (a) hydrogen;
- (b) halo especially fluoro, chloro or bromo.
- 5 (c) loweralkoxy especially C₁₋₆ alkoxy, e.g., methoxy, ethoxy, isopropoxy, t-butoxy or cyclohexyloxy;
- (d) lower alkylthio especially C₁₋₆ alkylthio, e.g., methylthio, ethylthio or
10 cyclohexylthio;
- (e) lower alkyl sulfinyl especially C₁₋₆ alkyl sulfinyl, e.g., methyl sulfinyl, i-propyl sulfinyl, and cyclopentyl sulfinyl.
- (f) lower alkyl sulfonyl especially C₁₋₆ alkyl
15 sulfonyl such as methyl sulfonyl, ethyl sulfonyl and n-butyl sulfonyl;
- (g) unsubstituted or substituted phenyl loweralkoxy such as benzyloxy;
- (h) loweralkyl especially C₁₋₆ alkyl such as
20 methyl, ethyl, propyl, t-butyl, pentyl, benzyl, cyclopropyl, cyclopentyl or cyclohexyl;
- (i) loweralkenyl especially C₂₋₆ alkenyl, for example, vinyl, allyl, and buten-2-yl.
- 25 (j) acetoxy;
- (k) -O-CH₂-O-;
- (l) -COOH;
- (m) aryl especially phenyl or substituted phenyl, e.g., 4-methoxyphenol,
30 2,4-difluorophenyl or 3-chlorophenyl; or
- (n) aryloxy especially phenoxy;

R¹ is

- (a) unsubstituted or substituted phenyl especially phenyl, 3-aminophenyl, 4-(3,3-dimethylbutyrylamino)phenyl, 3-methylphenyl, 3,5-dimethoxyphenyl, 3-ethylthiophenyl, 4-chlorophenyl, 4-trifluoromethylphenyl, 2,4-difluorophenyl, 4-nitrophenyl or the like;
- (b) unsubstituted or substituted heteroaryl especially monoheteroaryl such as thiadiazolyl, pyrrolyl, pyridyl or pyrazinyl, imidazolyl, thiaimidazolyl, quinolino or tetrazolyl or the like. The heteroaryl can be substituted with one or more of C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, halo especially chloro or fluoro, or the like; and

n is 1 to 5.

In a preferred embodiment of this invention,

R is

- (a) hydrogen;
(b) loweralkoxy;
(c) lower alkylthio;
(d) halo; or
(e) loweralkyl;


R¹ is unsubstituted or substituted phenyl; and
n is 1 to 3.

In the most preferred embodiment of this invention,

R is

- (a) hydrogen;
(b) C₁₋₃alkoxy such as methoxy, ethoxy or isopropoxy; or
(c) C₁₋₃ alkylthio such as methylthio, ethylthio or propylthio;

5 Step (1): treating an appropriately substituted
 2-hydroxy-benzaldehyde of the structural
 formula (II)



15 Step (2): treating the resulting Schiff-base with a
 reducing agent such as hydrogen in the
 presence of a catalyst, e.g., Pd/C or
 sodium tetrahydroboron.

In Step (1), the Schiff-base formation is well-documented. See M. M. Sprung, Chem. Rev., 26, 297 (1940); R. W. Layer, Chem. Rev., 63, 489 (1963); and G. Hilgetag and A. Martini, Preparative Organic Chemistry, Wiley-Interscience, pp. 504-508 (1968). Generally, the benzaldehyde is treated with neat amine or a solution thereof at relatively low temperatures, e.g., from about 0°C to about 25°C with adequate stirring until the reaction is substantially complete. The solution of the amine is usually prepared by dissolving the amine in an organic solvent preferably water, lower alcohol such as methanol, ethanol, isopropyl alcohol or an aqueous solution thereof. Under optimum conditions the reaction usually takes from about 1 to about 24 hours.

In most cases, Step (2) can be conveniently merged into Step (1) by the simple addition of a reducing agent to the reaction mixture of step (1). The crude Schiff-base in the reaction mixture is thus
5 reduced in situ to the desired hydroxybenzylamino derivatives of formula (I).

The most commonly used reducing agents, among others, are hydrogen in the presence of Pd/C and sodium borohydride. Other metal hydrides, for
10 example, lithium borohydride, lithium aluminumhydride NaCNBH_3 and other substituted aluminum or borohydrides may also be used. See H. O. House, Modern Synthetic Reactions, 2nd ed., W. A. Benjamin, Inc., 1972, pp. 45-54.

15 The reduction is usually conducted at relatively low temperatures, e.g., from about 0°C to about 30°C, preferably below 25°C. A solvent is usually required. The commonly used solvents for each reducing agent are summarized below in Table I.

20

TABLE 1

| Reducing Agent | Useful reaction solvents |
|--------------------------|-----------------------------------------------------------------------|
| 25 NaBH_4 | H_2O , CH_3OH , EtOH, i-PrOH, diglyme |
| LiBH_4 | Tetrahydrofuran, diglyme |
| 30 LiAlH_4 | Diethylether, tetrahydrofuran, 1,2-dimethoxyethane, and diglyme |
| $\text{H}_2/\text{Pd/C}$ | Ethylacetate, ethanol, water, acetic acid |

Usually the reaction is substantially complete within from about 1 hour to about 48 hours. However, under optimum conditions, the reaction may take only 1 to 6 hours.

5 The starting materials in the preparation of the novel hydroxybenzylamine derivatives are mostly commercially available or can be easily prepared via conventional methods, for example, 4-(3,3-dimethyl-
10 butyrylamino)aniline is simply derived from the reaction between 4-nitro-aniline and 3,3-dimethyl-butyryl chloride followed by hydrogenation.

As the novel compounds of this invention are organic bases, their pharmaceutically acceptable salts are those resulting from the neutralization of
15 the base with an acid. The acid employed is usually an inorganic acid such as a hydrohalic acid, e.g., hydrochloric or hydrobromic acid; sulfuric acid; nitric acid; or phosphoric acid. An organic acid
20 such as maleic, fumaric, tartaric, citric, acetic, salicylic, succinic, benzoic, benzenesulfonic, glutamic, lactic or isethionic acid is also commonly used. Generally the neutralization is conducted in
25 an inert solvent such as water; a C₁₋₃ alkanol such as methanol, ethanol or isopropanol; a C₃₋₆-ketone such as acetone, or ethylmethyl ketone; an ethereal solvent such as diethyl ether, tetrahydrofuran or dioxane; acetonitrile; or an aromatic solvent such as toluene. Mixtures of the above described solvents
30 are also employed. Generally the neutralization is carried out in aqueous ethanol, at 0°-75°C,

preferably at 0°-25°C, followed by filtration to collect the salts.

This invention also relates to a method of treating topical inflammation in patients in need of such treatment. Generally, a sufficient amount of a compound of formula (I) or a pharmaceutical composition thereof, particularly an especially preferred compound, is administered to the patient as the active constituent.

The U.V. erythema assay by which topical anti-inflammatory activity is determined, is based on the ability of the compounds of Formula I to inhibit the erythema induced on the skin of guinea pigs by UV radiation. It has been substantiated by recent studies that prostaglandin synthesis and the release thereof may be the primary process involved in the production of erythema. A protocol of the assay and some results thereof are summarized below in table II.

TABLE II

U.V. Erythema Assay

Procedure

Guinea pigs were depilated and stabilized for 30 minutes. Following anesthesia with Nembutal (35 mg/kg), they were exposed to a U.V. lamp for 45 minutes at a distance of seven inches from the abdominal surface. Compound was applied topically (seven applications in a total of 0.1 ml vehicle) 15 minutes post exposure. The surface was then gently washed and read two hours post treatment.

Vehicle 1774 - 85% EtOH, 12% Propylene Glycol, 3%
Methyl Salicylate

Erythema Scoring

| | | | |
|----|--------------------|--------------|---------------------------------|
| 5 | Numerical | | |
| | <u>Designation</u> | <u>Score</u> | <u>Description</u> |
| | Negative | 0 | No erythema (normal skin color) |
| | Trace | + 1 | Faint pink in some areas |
| 10 | Slight | + 2 | Faint pink over entire site |
| | Slight- | | |
| | Moderate | + 3 | Faint pink to red |
| | Moderate | + 4 | Red |
| | Marked | + 5 | Red to purplish red |

15

Evaluation of Erythema

A numerical value 0-5 was assigned to the degree of erythema observed under standard lighting conditions. Drug effect on the developing erythema was calculated as follows:

20

$$\% \text{ suppression} = \frac{\left(\frac{\text{score of vehicle}}{\text{treated}} \right) - \left(\frac{\text{Score of drug}}{\text{Treated}} \right)}{\text{score of vehicle treated}} \times 100$$

25

Results: % Inhibition of Erythema by N-(2-hydroxy-5-methoxybenzyl)aniline

30

0081782

- 11 -

16654

| Dosage (mg) | % Inhibition (mean of 3 animals) | | | | |
|----------------|----------------------------------|-------|-------|-------|--------|
| | 2 Hr. | 3 Hr. | 4 Hr. | 6 Hr. | 24 Hr. |
| 0.1 | 19.3 | 16.7 | 15.0 | 6.7 | 0 |
| 5 0.3 | 55.3 | 50.0 | 46.7 | 35.0 | 0 |
| 1.0 | 80.3 | 91.7 | 76.7 | 63.3 | 26.7 |
| 3.0 | 100 | 100 | 63.3 | 43.3 | 33.3 |

For treatment of inflammation, fever or
10 pain, the compounds of the invention are administered
topically, by inhalation spray or rectally in dosage
unit formulations containing conventional non-toxic
pharmaceutically acceptable carriers, adjuvants and
vehicles. In addition to the treatment of
15 warm-blooded animals such as mice, rats, horses,
dogs, cats, etc., the compounds of the invention are
effective in the treatment of humans.

The pharmaceutical compositions containing
the active ingredient may be in a form suitable for
20 topical use, for example, aqueous or oily solutions
or suspensions, dispersible powders or granules,
tinctures, topical aerosol emulsions, creams,
ointments, jellies, suppositories or the like.
Compositions intended for topical use may be prepared
25 according to any method known to the art for the
manufacture of pharmaceutical compositions
and such compositions may contain one or more active
compounds.

Aqueous suspensions contain the active
30 materials in admixture with excipients suitable for
the manufacture of aqueous suspensions. Such

excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyoxyethylene sorbitan monooleate. The said aqueous suspensions may also contain one or more preservatives, for example, ethyl, or n-propyl p-hydroxybenzoate.

Oily suspension may be formulated by suspending the active ingredient in a vegetable oil, for example, arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example, beeswax, hard paraffin or cetyl alcohol. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the

addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oils, or a mineral oil, for example, liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example, gum acacia or gum tragacanth, naturally-occurring phosphatides, for example, soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example, sorbitan mono-oleate, and condensation products of the said partial esters with ethylene oxide, for example, polyoxyethylene sorbitan monooleate.

An ointment containing the pharmaceutical compositions of the present invention may be prepared, among other methods known in the art, by combining the active ingredient with a medium consisting of a glycol, a lower alkanol, and water; a gelling agent; and optionally an adjuvant such as diisopropyl adipate, diethyl sebacate, ethyl carproate and ethyl laurate. Suitable glycols include propylene glycol, butylene glycol, polyethylene glycol and the like. Generally, a

carboxyvinyl polymer preneutralized with an organic amine such as diisopropyl amine and triethylamine, or a cellulose, e.g., hydroxyethyl cellulose, methyl cellulose, carboxymethyl cellulose, hydroxypropyl cellulose, is used as the gelling agent.

The compounds of the invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Dosage levels of the order to 0.2 mg to 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (10 mg to 7 gms per patient per day). For example, inflammation is effectively treated by the administration from about 0.5 to 50 mg of the compound per kilogram of body weight per day (25 mg to 5 gms per patient per day). Advantageously, from about 2 mg to about 20 mg per kilogram of body weight per daily dosage produces highly effective results (50 mg to 1 gm per patient per day).

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from

about 25 mg to about 1 g of active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

EXAMPLE 1

4-(2-hydroxy-5-methoxybenzlamino)-3,3-dimethylbutyryl-aminobenzene

Step A: Preparation of (3,3-dimethylbutyryl-amino)aniline

Five grams of 4-nitroaniline was stirred in 25 ml of pyridine and 4.95 g of 3,3-dimethylbutyryl chloride was added. The mixture was stirred 15 hours and poured into ice-water. The resultant solid was collected on a filter and dissolved in 200 ml of ethyl acetate and the column was treated with activated charcoal and concentrated to yield 6.3 g of 4-nitro-3,3-dimethylbutyrylaminobenzene, m.p. 192-194°. This material was reduced in 200 ml of methanol, at 3 atmospheres of hydrogen in the presence of 5% Pd/carbon. After removal of the catalyst by filtration, concentration of the filtrate gives 5.3 g of product (3,3-dimethylbutyrylamino)-aniline, m.p. 138-140°C.

Step B: Preparation of 4-(2-hydroxy-5-methoxybenzyl-amino)-3,3-dimethylbutyrylamino benzene

A mixture of 2.5 g of 3,3-dimethylbutyryl-aminoaniline and 1.84 g of 2-hydroxy-5-methoxybenzal-
5 dehyde was stirred in 50 ml of methanol, under a nitrogen atmosphere, for 15 hours. The mixture was then refluxed until tlc indicates that most of the starting material had reacted. After cooling, 0.50 g of NaBH_4 was added in two batches, and the mixture
10 was stirred 2.5 hours. After concentration 6.9 g of solid was collected which was triturated with 100 ml of water and enough 2.5N HCl to neutralize the supernatant. 3.95 g of crude product was collected and purified by treatment with activated charcoal and
15 recrystallization from ethyl acetate to give 2.5 g of 4-(2-hydroxy-5-methoxybenzylamino)-3,3-dimethylbutyrylamino benzene, m.p. 126-129°C.

Following substantially the same procedure as described above but substituting for the starting
20 materials used therein 5-(t-butyl)-2-hydroxy-3-iodobenzaldehyde and 4-acetaminoaniline, there was prepared 4-(5-t-butyl-2-hydroxy-3-iodobenzylamino)acetanilide. M.p. 134.5-136°C.

25 EXAMPLE 2
3-(3,5-Dichloro-2-Hydroxybenzylamino)[1,2,5]thia-
diazole

A mixture of 6.9 g of 3,5-dichloro-2-hydroxybenzaldehyde and 3-aminó[1,2,5]thiazole hydro-
30 chloride was heated at 80° for 18 hours. After

cooling, 6.9 g of precipitate, m.p. 174-175° was collected. This precipitate was added to 100 ml of methanol and the mixture was cooled in an ice-bath before 1.4 g of NaBH_4 is added. The mixture was stirred 1/2 hour and concentrated to give 9.0 g of solid which was purified by trituration with water and enough 2.5N HCl to give a neutral supernatant. The insolubles were then collected, heated with charcoal and recrystallized from methylene chloride to give 5.1 g of 3-(3,5-dichloro-2-hydroxybenzyl-amino)[1,2,5]thiadiazole. M.p. 131-132°C.

EXAMPLE 3

N-(2-Hydroxybenzyl)Aniline

A mixture of 6.1 g of salicylaldehyde, 4.7 g of aniline and 50 ml of methanol was stirred and heated at 80° for 17 hours, cooled to 25° and 2.0 g of sodium borohydride was added in two batches. The mixture was stirred for 1 hour and the resultant precipitate was collected on a filter. This solid was treated with activated charcoal and recrystallized from 150 ml of isopropanol to give 5.4 g of N-(2-hydroxybenzyl)aniline. M.p. 110-111°C.

EXAMPLE 4

N-(2-Hydroxy-5-Methoxybenzyl)Aniline

A mixture of 15.2 g 2-hydroxy-5-methoxy benzaldehyde, 93 g of aniline and 100 ml of methanol was stirred at 25° for 20 hours. The mixture was then cooled with an ice-bath and stirred, and 3.9 g

of sodium borohydride was added slowly. The mixture was stirred for 1 hour at 25°, 150 ml of water was added and the pH was adjusted to 7.0 with 2.5N hydrochloric acid. The resultant solid was
5 collected, washed with water and purified by chromatography on silica gel using 3:17 ethyl acetate: n-hexane as an eluant. A 17.7 g yield of N-(2-hydroxy-5-methoxybenzyl)aniline was obtained. M.p. 82-84°.

10

EXAMPLE 5

N-(2-Hydroxy-5-methoxybenzyl)Aniline

A mixture of 15.2 g of 2-hydroxy-5-methoxybenzaldehyde, 15 g of aniline and 200 ml of methanol
15 was stirred for 3 hours. The mixture was then hydrogenated at 40 psi in the presence of 2.0 g of 10% Pd/C until 0.1 mole of hydrogen was consumed. The catalyst was removed by filtration and the filtrate was concentrated in vacuo to a yellow oil which was
20 crystallized from ether n-hexane to yield 18 g of N-(2-hydroxy-5-methoxybenzyl)aniline. M.p. 82.5- 83°C.

Following substantially the same procedure as described above, but substituting for aniline used
25 therein the following compounds:

- (1) 2,6-dichloroaniline
- (2) 3,5-dimethylaniline
- (3) 3,5-dimethoxyaniline
- (4) 4-trifluoromethylaniline

30

There are obtained the following
corresponding hydroxybenzylaminobenzene derivatives:

- (1) 2,6-dichloro-N-(2-hydroxy-5-methoxybenzyl)-
aniline, m.p.
- 5 (2) 3,5-dimethyl-N-(2-hydroxy-5-methoxybenzyl)-
aniline
- (3) 3,5-dimethoxy-N-(2-hydroxy-5-methoxybenzyl)-
aniline
- 10 (4) 4-trifluoromethylaniline

EXAMPLE 6

Following substantially the same procedure
as described in Example 5, but substituting for
aniline used therein the following heterocyclic
15 amines:

- (1) 3-aminoquinoline
- (2) 2-amino-4,6-dimethylpyridine

There were obtained the corresponding hydroxybenzyl
amine derivatives as listed below:

- 20 (1) 3-[N-(2-hydroxy-5-methoxybenzyl)amino]-
quinoline, m.p. 101-103°C.
- (2) 2-[N-(2-hydroxy-5-methoxybenzyl)amino]-4,6-
dimethylpyridine m.p. 95-96°C.

EXAMPLE 7

Set forth below are some illustrative topical
formulations containing a selected active compound of
the instant invention.

Formulation Number 1 - Solution

3-(3,5-dichloro-2-hydroxybenzylamino) [1,2,5]
thiadiazole (a) - 2.5%
Distilled water qs to 100%

5 Procedure: Dissolve compound (a) in enough
water to make 100%. Filter the solution. Apply to
the affected area.

Formulation Number 2 - Tincture

N-(2-hydroxybenzyl)aniline (b) - 2.5%
10 Alcohol U.S.P. - 50%
Water qs to 100%

Procedure: Dissolve compound (b) in the
alcohol. Add sufficient water to make 100%. Filter
and apply to affected area.

15

Formulation Number 3 - Topical Aerosol

N-(2-hydroxy-5-methoxy-
benzyl)Aniline (c) - 2.5%
Alcohol U.S.P. - 5%
20 Isopropylmyristate - 5%
Conventional halogenated hydrocarbon propellant qs
100% e.g., Freon 11(trichlorofluoromethane), Freon
12(dichlorodifluoromethane), Freon 14 (carbon
tetrafluoride), Freon C 318 (Octafluorocyclobutane),
25 Freon 114(Cryofluorane), etc.

Procedure: Dissolve Compound (c) in the
alcohol and isopropylmyristate. Add sufficient
halogenated propellant and introduce into
conventional aerosol containers either by pressure or
30 by cold filing. Apply to affected area.

0081782

- 21 -

16654

Formulation Number 4 - Ointment

4-(2-hydroxy-5-methoxybenzylamino)-3,3-di-
methylbutyrylamino benzene (d) - 2.5%

Petrolatum U.S.P. qs to 100%

5 Procedure: Heat the petrolatum to 60°C.
Add compound (d) and stir until thoroughly
dispersed. Cool to room temperature. Apply to
affected area.

10

15


20

25

30

WHAT IS CLAIMED IS:

1. A compound of structural formula (I)



(I)

10 wherein:

R is (a) fluoro;

(b) methoxy, ethoxy, n-propoxy or i-propoxy;

(c) methylthio, ethylthio, n-propylthio or
i-propylthio;

15 (d) -OCH₂-O;

(e) -COOH; or

(f) aryloxy;

R₁ is (a) unsubstituted or substituted phenyl; or


(b) unsubstituted or substituted heteroaryl;

20 n is 1 or 2.

2. The compound of Claim 1 wherein the
compound is N-(2-hydroxy-5-methoxybenzyl)aniline.

25 3. A process for the preparation of a
compound of structural formula (I) according to Claim
1 comprising:

(a) treating an appropriately substituted
2-hydroxybenzaldehyde of the
30 structural formula II



(II)

10 with an amine of formula R^1NH_2 to form a Schiff-base; and
(b) treating the Schiff-base with a reducing agent.

4. The process of Claim 3 for preparing
15 N-(2-hydroxy-5-methoxybenzyl)aniline.

5. A pharmaceutical composition for treatment of topical inflammation comprising a
20 pharmaceutical carrier and an effective amount of a compound of formula (I) according to Claim 1.

6. The pharmaceutical composition of Claim 7 wherein the compound is N-(2-hydroxy-5-methoxy-
25 benzyl)aniline.

30


35

1

CLAIMS FOR AUSTRIA

1. A process for the preparation of a compound of structural formula (I)

5



10

(I)

wherein:

R is (a) fluoro;

(b) methoxy, ethoxy, n-propoxy or i-propoxy;

(c) methylthio, ethylthio, n-propylthio or i-propylthio;

15

(d) -OCH₂-O;

(e) -COOH; or

(f) aryloxy;

R¹ is (a) unsubstituted or substituted phenyl; or

20


(b) unsubstituted or substituted heteroaryl;

n is 1 or 2

comprising:

(a) treating an appropriately substituted 2-hydroxybenzaldehyde of the structural formula II

25



30

(II)

with an amine of formula R¹NH₂ to form a Schiff-base; and

35

(b) treating the Schiff-base with a reducing agent.

1

16654

2. The process of Claim 1 for preparing
N-(2-hydroxy-5-methoxybenzyl)aniline.

5

10

15

20

25

30

35



| DOCUMENTS CONSIDERED TO BE RELEVANT | | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Category | Citation of document with indication, where appropriate, of relevant passages | Relevant to claim | CLASSIFICATION OF THE APPLICATION (Int. Cl. 3) |
| X | CHEMICAL ABSTRACTS, vol. 86, no. 9, 28th February 1977, page 366, no. 54830q, Columbus Ohio (USA); J.STRADINS et al.: "Comparative voltammetric study of para-substituted phenols and their N-phenylaminomethyl derivatives on a graphite anode". & ZH. ORG. KHIM. 1976, 12(9), 1949-55. *Abstract* | 1-4 | C 07 C 93/14 C 07 C 91/30 C 07 C 103/44 C 07 D 285/10 C 07 D 213/74 C 07 D 215/38 A 61 K 31/135 A 61 K 31/41 A 61 K 31/445 A 61 K 31/47 |
| X | --- ARCHIV DER PHARMAZIE, vol. 298, no. 7, 1965, pages 423-434, Weinheim (DE); R.POHLUDEK-FABINI et al.: "Verhalten rhodanierter Schiffscher Basen gegenüber Raney-Nickel, Hydrazin und Alkalien". *Page 429* | 1-4 | |
| A | --- US-A-2 784 138 (R.WEGLER) *Column 1, lines 15-28* | 1 | |
| A | --- US-A-3 996 278 (Z.H.SCHLAGER) *Column 1, lines 21-43; column 2, line 62 - column 3, line 2* | 1 | |
| | ----- | | |
| The present search report has been drawn up for all claims | | | |
| Place of search THE HAGUE | | Date of completion of the search 18-03-1983 | Examiner PAUWELS G.R.A. |
| CATEGORY OF CITED DOCUMENTS | | | |
| X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document | | T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document | |